

Rabbit anti-Human CDA Polyclonal Antibody | anti-CDA antibody

Anti-CDA Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257620	Rabbit	Human	Immunohistochemistry, Western Blot

PRODUCT SYNONYMS


CDA; Polyclonal Antibody; Anti-CDA Antibody; Rabbit Polyclonal; CDA Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CDD Antibody; cytidine deaminase; anti-CDA antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CDA
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CDA antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human CDA fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human CDA fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IHC (Immunohistochemistry)
Immunochemical staining of human CDA in human spleen with rabbit polyclonal antibody at 1:5000 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)
Anti-CDA rabbit polyclonal antibody at 1:500 dilutionLane A: HeLa Whole Cell LysateLane B: U-251 MG Whole Cell LysateLysates/proteins at 30 ug per lane.SecondaryGoat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.Developed using the ECL technique.Performed under reducing conditions.Predicted band size:16 kDaObserved band size:16 kDa(We are unsure as to the identity of these extra bands.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cda-antibody-immunohistochemistry-western-blot-elisa/257620>