

Rabbit anti-Human ASPSCR1 Polyclonal Antibody | anti-ASPSCR1 antibody

Anti-ASPSCR1 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257654	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ASPSCR1; Polyclonal Antibody; Anti-ASPSCR1 Antibody; Rabbit Polyclonal; ASPSCR1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-ASPSCR1 Antibody; Anti-ASPL Antibody; Anti-ASPS Antibody; Anti-RCC17 Antibody; Anti-TUG Antibody; Anti-UBXD9 Antibody; Anti-UBXN9 Antibody; anti-ASPSCR1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human ASPSCR1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-ASPSCR1 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human ASPSCR1 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human ASPSCR1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IP (Immunoprecipitation)

ASPSCR1 was immunoprecipitated using: Lane A: 0.5 mg K562 Whole Cell Lysate, 4 uL anti-ASPSCR1 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-ASPSCR1 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 60 kDa. Observed band size: ~72 kDa.




IF (Immunofluorescence)


Immunofluorescence staining of ASPSCR1 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human ASPSCR1 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus, Cytoplasm and cell membrane.

**IHC (Immunohistochemisry)**

Immunochemical staining of human ASPSCR1 in human liver with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohiostchemistry)**

Immunochemical staining of human ASPSCR1 in human kidney with rabbit polyclonal antibody at 1:1000 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-ASPSCR1 rabbit polyclonal antibody at 1:500 dilutionLane A: K562 Whole Cell LysateLane B: 293T Whole Cell LysateLane C: SW480 Whole Cell LysateLysates/proteins at 30 ug per lane.SecondaryGoat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.Developed using the ECL technique.Performed under reducing conditions.Predicted band size:60 kDaObserved band size:72 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/aspscr1-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/257654>