

Rabbit anti-Human APBB3 Polyclonal Antibody | anti-APBB3 antibody

Anti-APBB3 Antibody, Rabbit Polyclonal


CATALOG # AAA257693	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

APBB3; Polyclonal Antibody; Anti-APBB3 Antibody; Rabbit Polyclonal; APBB3 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-FE65L2 Antibody; Anti-SRA Antibody; anti-APBB3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human APBB3
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-APBB3 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E Coli-derived Human APBB3 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human APBB3 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IP (Immunoprecipitation)

APBB3 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate, 4 uL anti-APBB3 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-APBB3 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 53 kDa. Observed band size: 53 kDa.



WB (Western Blot)

Anti-APBB3 rabbit polyclonal antibody at 1:500 dilution. Lane A: HeLa Whole Cell Lysate. Lane B: A431 Whole Cell Lysate. Lane C: 293T Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 53 kDa. Observed band size: 53 kDa. (We are unsure as to the identity of these extra bands.)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.