

Rabbit anti-Human TERF2IP Polyclonal Antibody | anti-TERF2IP antibody

Anti-TERF2IP Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257702	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

TERF2IP; Polyclonal Antibody; Anti-TERF2IP Antibody; Rabbit Polyclonal; TERF2IP Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-DRIP5 Antibody; Anti-RAP1 Antibody; anti-TERF2IP antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human TERF2IP
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-TERF2IP antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human TERF2IP fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human TERF2IP fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.



IF (Immunofluorescence)

Immunofluorescence staining of TERF2IP in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human TERF2IP polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



WB (Western Blot)

Anti-TERF2IP rabbit polyclonal antibody at 1:500 dilution
Lane A: HeLa Whole Cell Lysate
Lane B: HEK293 Whole Cell Lysate
Lane C: HepG2 Whole Cell Lysate
Lane D: A431 Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 36 kDa Observed band size: 60 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.