

Rabbit anti-Human SEPT9 Polyclonal Antibody | anti-SEPT9 antibody

Anti-SEPT9 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257731	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

SEPT9; Polyclonal Antibody; Anti-SEPT9 Antibody; Rabbit Polyclonal; SEPT9 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-AF17q25 Antibody; Anti-MSF Antibody; Anti-MSF1 Antibody; Anti-NAPB Antibody; Anti-PNUTL4 Antibody; Anti-SeptD1 Antibody; Anti-SINT1 Antibody; anti-SEPT9 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human SEPT9
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-SEPT9 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human SEPT9 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human SEPT9 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	<p>This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.</p> <p>This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.</p>

IMAGES



IP (Immunoprecipitation)

SEPT9 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate, 4 uL anti-SEPT9 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-SEPT9 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 65 kDa. Observed band size: 65 kDa.



IF (Immunofluorescence)


Immunofluorescence staining of SEPT9 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human SEPT9 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to.

**IHC (Immunohistochemistry)**

Immunochemical staining of human SEPT9 in human lymph node with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human SEPT9 in human testis with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-SEPT9 rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 65 kDa Observed band size: 72 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sept9-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/257731>