

Rabbit anti-Human DcR1/TRAILR3 Polyclonal Antibody | anti-TNFRSF10C antibody

Anti-DcR1/TRAILR3 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257751	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

DcR1/TRAILR3; Polyclonal Antibody; Anti-DcR1/TRAILR3 Antibody; Rabbit Polyclonal; DcR1/TRAILR3 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CD263 Antibody; Anti-DCR1 Antibody; Anti-DCR1-TNFR Antibody; Anti-LIT Antibody; Anti-TRAIL-R3 Antibody; Anti-TRAILR3 Antibody; Anti-TRID Antibody; tumor necrosis factor receptor superfamily member 10c; anti-TNFRSF10C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human DcR1/TRAILR3
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-TNFRSF10C antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human DcR1/TRAILR3 fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human DcR1/TRAILR3 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES



IP (Immunoprecipitation)

TNFRSF10C was immunoprecipitated using: Lane A: 0.5 mg Hela Whole Cell Lysate 4 uL anti-TNFRSF10C rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-TNFRSF10C rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 65 kDa. Observed band size: 65 kDa.



IF (Immunofluorescence)

Immunofluorescence staining of TRAILR3 in A549 cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human TRAILR3 polyclonal antibody (dilution ratio 1:200) at 4 degree C overnight. Then cells were stained with the Alexa Fluor-488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



WB (Western Blot)

Anti-DcR1/TRAILR3 rabbit polyclonal antibody at 1:500 dilutionLane A: HeLa Whole Cell LysateLane B: Jurkat Whole Cell LysateLane C: A549 Whole Cell LysateLysates/proteins at 30 ug per lane.SecondaryGoat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.Developed using the ECL technique.Performed under reducing conditions.Predicted band size:65 kDaObserved band size:70 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TNFRSF10C antibody	Background: TNFRSF10C CNV in patients with CRC is associated with distant metastatic disease. A high frequency of CGI methylation in the TNFRSF10C promoter results in inactivation of the gene and enhancement of tumor growth in most PC cell lines (except CFPAC-1). Inactivation of TNFRSF10C by CpG island (CGI) hypermethylation can play an important role in PC progression and be potentially useful as a test marker and a new therapeutic approach for PC.
Product Categories/Family for anti-TNFRSF10C antibody	Death Receptor Signaling

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/dcr1-trailr3-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-western-blot-elisa/257751>