

Rabbit anti-Human AIFM1/AIF Polyclonal Antibody | anti-AIFM1 antibody

Anti-AIFM1/AIF Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA257761	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

AIFM1/AIF; Polyclonal Antibody; Anti-AIFM1/AIF Antibody; Rabbit Polyclonal; AIFM1/AIF Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-AIF Antibody; Anti-CMT2D Antibody; Anti-CMTX4 Antibody; Anti-COWCK Antibody; Anti-COXP6 Antibody; Anti-NADMR Antibody; Anti-NAMSD Antibody; Anti-PDCD8 Antibody; apoptosis inducing factor; mitochondria associated 1; anti-AIFM1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human AIFM1/AIF
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid. PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-AIFM1 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E Coli-derived Human AIFM1/AIF fragment
Conjugation	Unconjugated
Preparation	Produced in rabbits immunized with E Coli-derived Human AIFM1/AIF fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.




IP (Immunoprecipitation)

AIFM1 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate 4 uL anti-AIFM1 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-AIFM1 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 67 kDa Observed band size : 67 kDa




IHC (Immunohistochemistry)

Immunochemical staining of human AIFM1 in human liver with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human AIFM1 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-AIFM1/AIF rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lane B: HepG2 Whole Cell Lysate Lane C: K562 Whole Cell Lysate Lane D: MCF7 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 67 kDa Observed band size: 67 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.