

PD-L2 Active Protein | PDCD1LG2 active protein

PD-L2 Protein, Human, Recombinant (ECD, His Tag)

CATALOG #
AAA257866

HOST
HEK293 Cells



PRODUCT SYNONYMS

PD-L2; PD-L2 Protein; Human; Recombinant (ECD; His Tag); Human B7-DC/PD-L2/CD273 Protein (His Tag); B7-DC Protein; B7DC Protein; bA574F11.2 Protein; Btdc Protein; CD273 Protein; PDCD1L2 Protein; PDL2 Protein; programmed cell death 1 ligand 2; PDCD1LG2 active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>98% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Pro219
Species	Human
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Leu 20
Tag	C-His
Protein Construction	A DNA sequence encoding the extracellular domain (Met 1-Pro 219) of human PD-L2 (NP_079515.2) was expressed with a C-terminal polyhistidine tag.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Activity	Measured by its binding ability in a functional ELISA. Immobilized human PD-L2 at 2ug/ml (100uL/well) can bind human PD-1. The EC50 of human PD-1 is 150-800ng/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.

IMAGES



Application Data

Measured by its binding ability in a functional ELISA. Immobilized human PD-L2 at 2 ug/ml (100 ul/well) can bind human PD-1, the EC50 of human PD-1 is 150-800 ng/mL.

SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for PDCD1LG2 active protein	Background: Programmed death ligand 2 (PD-L2), also referred to as B7-DC and CD273, is a member of the B7 family of proteins including B7-1, B7-2, B7-H2, B7-H1 (PD-L1), and B7-H3. PD-L2 is a type I membrane protein and structurally consists of an extracellular region containing one V-like and one C-like Ig domain, a transmembrane region, and a short cytoplasmic domain. PD-L2 is expressed on antigen presenting cells, placental endothelium and medullary thymic epithelial cells, and can be induced by LPS in B cells, INF-? in monocytes, or LPS plus IFN-? in dendritic cells. The CD28 and B7 protein families are critical regulators of immune responses. PD-L2 and PD-L1 are two ligands for PD-1, member of the CD28/CTLA4 family expressed on activated lymphoid cells, and thus provide signals for regulating T cell activation and immune tolerance. The interaction of B7-DC/PD-1 exhibited a 2-6-fold higher affinity compared with the interaction of B7-H1/PD-1.
Product Categories/Family for PDCD1LG2 active protein	Cancer Drug Targets
References	Latchman Y, et al. (2001) PD-L2 is a second ligand for PD-1 and inhibits T cell activation. Nat Immunol. 2: 261-8.Carreno BM, et al. (2005) Therapeutic opportunities in the B7/CD28 family of ligands and receptors. Curr Opin Pharmacol. 5(4): 424-30.Radhakrishnan S, et al. (2007) B7-DC/PD-L2 cross-linking induces NF-kappaB-dependent protection of dendritic cells from cell death. J Immunol. 178(3): 1426-32.

NCBI AND UNIPROT INFORMATION

NCBI GI #	190014605
NCBI GeneID	80380
NCBI Accession #	NP_079515.2
NCBI GenBank Nucleotide #	NM_025239.3
UniProt Accession #	Q9BQ51; Q14CN8; Q5T7Z6; Q6JXL8; Q6JXL9
Molecular Weight	20,745 Da
NCBI Official Full Name	programmed cell death 1 ligand 2
NCBI Official Synonym Full Names	programmed cell death 1 ligand 2
NCBI Official Symbol	PDCD1LG2
NCBI Official Synonym Symbols	B7DC; Btdc; PDL2; CD273; PD-L2; PDCD1L2; bA574F11.2
NCBI Protein Information	programmed cell death 1 ligand 2
UniProt Protein Name	Programmed cell death 1 ligand 2
UniProt Gene Name	PDCD1LG2

UniProt Synonym Gene Names	B7DC; CD273; PDCD1L2; PDL2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; B7-DC
UniProt Entry Name	PD1L2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-pcd1lg2-active-protein/pd-l2/257866>