

FGFR4 active protein

FGFR4 Protein, Human, Recombinant (His Tag)

CATALOG #
AAA257892


HOST
HEK293 Cells

PRODUCT SYNONYMS

FGFR4; FGFR4 Protein; Human; Recombinant (His Tag); Human FGFR4/FGF Receptor 4 Protein (His Tag); CD334 Protein; JTK2 Protein; TKF Protein; fibroblast growth factor receptor 4; FGFR4 active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>90% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Asp36 9
Species	Human
Activity	Measured by its ability to inhibit FGF acidic (aFGF/FGF1) dependent proliferation of Balb/c3T3 mouse embryonic fibroblasts. The ED50 for this effect is typically 0.2-1ug/mL.
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Leu 22
Tag	C-His
Protein Construction	A DNA sequence encoding the extracellular domain (Met 1-Asp 369) of human FGFR4 (NP_002002.3) was fused with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data

Measured by its ability to inhibit FGF acidic (aFGF / FGF1) dependent proliferation of Balb/c3T3 mouse embryonic fibroblasts. The ED50 for this effect is typically 0.2-1ug/mL.

ADDITIONAL INFORMATION

Related Product Information for FGFR4 active protein	Background: Fibroblast growth factor receptor 4 (FGFR4) also known as CD334 antigen or tyrosine kinase related to fibroblast growth factor receptor, is a member of the fibroblast growth factor receptor family, where amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of FGFR4/CD334 interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. FGFR4/CD334 preferentially binds acidic fibroblast growth factor and, although its specific function is unknown, it is overexpressed in gynecological tumor samples, suggesting a role in breast and ovarian tumorigenesis. FGFR4/CD334 signaling is down-regulated by receptor internalization and degradation; MMP14 promotes internalization and degradation of FGFR4/CD334. Mutations in FGFR4/CD334 lead to constitutive kinase activation or impair normal FGFR4 inactivation lead to aberrant signaling.
Product Categories/Family for FGFR4 active protein	Cancer Drug Targets Receptor Tyrosine Kinases (RTKs)
References	Hart KC, et al. (2000) Transformation and Stat activation by derivatives of FGFR1, FGFR3, and FGFR4. Oncogene. 19(29): 3309-20. Xie MH, et al. (1999) FGF-19, a novel fibroblast growth factor with unique specificity for FGFR4. Cytokine. 11(10): 729-35. Yu C, et al. (2000) Elevated cholesterol metabolism and bile acid synthesis in mice lacking membrane tyrosine kinase receptor FGFR4. J Biol Chem. 275(20): 15482-9.

NCBI AND UNIPROT INFORMATION

NCBI GI #	47524173
NCBI GeneID	2264
NCBI Accession #	NP_002002.3
NCBI GenBank Nucleotide #	NM_002011.3
UniProt Accession #	P22455; O43785; Q14309; Q71TW8; Q8TDA0; AAB25788
Molecular Weight	87,954 Da
NCBI Official Full Name	fibroblast growth factor receptor 4 isoform 1
NCBI Official Synonym Full Names	fibroblast growth factor receptor 4
NCBI Official Symbol	FGFR4
NCBI Official Synonym Symbols	TKF; JTK2; CD334

NCBI Protein Information	fibroblast growth factor receptor 4; FGFR-4; tyrosylprotein kinase; protein-tyrosine kinase; hydroxyaryl-protein kinase; tyrosine kinase related to fibroblast growth factor receptor
UniProt Protein Name	Fibroblast growth factor receptor 4
UniProt Gene Name	FGFR4
UniProt Synonym Gene Names	JTK2; TKF; FGFR-4
UniProt Entry Name	FGFR4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-active-protein/fgr4/257892>