

BTLA active protein

BTLA Protein, Human, Recombinant (hFc Tag)

CATALOG #
AAA257998

HOST
HEK293 Cells



PRODUCT SYNONYMS

BTLA; BTLA Protein; Human; Recombinant (hFc Tag); Human BTLA Protein (Fc Tag); BTLA1 Protein; CD272 Protein; B and T lymphocyte associated; BTLA active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>90% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Thr134
Species	Human
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Lys 31
Tag	C-human IgG1-Fc
Protein Construction	A DNA sequence encoding the human BTLA (NP_001078826.1)(Met1-Thr134) was expressed with the Fc region of human IgG1 at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Activity	1.Measured by its binding ability in a functional ELISA. 2.Immobilized human BTLA-Fc at 10ug/mL (100uL/well) can bind biotinylated mouse HVEM-Fch. The EC50 of biotinylated mouse HVEM-Fch is 6-70ng/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.

IMAGES



Application Data

Measured by its binding ability in a functional ELISA. Immobilized human BTLA-Fc at 10ug/mL (100uL/well) can bind biotinylated mouse HVEM-Fch, the EC50 of biotinylated mouse HVEM-Fch is 6-70 ng/mL.

SDS-PAGE

ADDITIONAL INFORMATION

Related Product Information for BTLA active protein	Background: BTLA is a inhibitory molecule which belongs to the Ig superfamily. It down-modulates immune responses. As such, reagents that regulate the binding of BTLA to its ligand or alter BTLA signaling have significant therapeutic promise. BTLA is crucial to understand the mechanism(s) of action of these antibodies before attempting clinical applications. BTLA is not expressed by naive T cells, but it is induced during activation and remains expressed on T helper type 1 (T(H)1) but not T(H)2 cells. BTLA is a third inhibitory receptor on T lymphocytes with similarities to cytotoxic T lymphocyte-associated antigen 4 (CTLA-4) and programmed death 1 (PD-1).
Product Categories/Family for BTLA active protein	Cancer Drug Targets ITIM/ITAM Immunoreceptors and Related Molecules
References	Fourcade J, et al. (2012) CD8(+) T cells specific for tumor antigens can be rendered dysfunctional by the tumor microenvironment through upregulation of the inhibitory receptors BTLA and PD-1. Cancer Res. 72(4):887-96.Kojima R, et al. (2011) Molecular basis for herpesvirus entry mediator recognition by the human immune inhibitory receptor CD160 and its relationship to the cosignaling molecules BTLA and LIGHT. J Mol Biol. 413(4):762-72.Oki M, et al. (2011) A functional polymorphism in B and T lymphocyte attenuator is associated with susceptibility to rheumatoid arthritis. Clin Dev Immunol. 305656.

NCBI AND UNIPROT INFORMATION

NCBI GI #	145580619
NCBI GeneID	151888
NCBI Accession #	NP_001078826.1
NCBI GenBank Nucleotide #	NM_001085357.1
UniProt Accession #	Q7Z6A9; Q3B831; Q3HS85; Q6ZNH9
Molecular Weight	32,834 Da
NCBI Official Full Name	B- and T-lymphocyte attenuator isoform 2
NCBI Official Synonym Full Names	B and T lymphocyte associated
NCBI Official Symbol	BTLA
NCBI Official Synonym Symbols	BTLA1; CD272
NCBI Protein Information	B- and T-lymphocyte attenuator; B and T lymphocyte attenuator; B- and T-lymphocyte-associated protein
UniProt Protein Name	B- and T-lymphocyte attenuator

UniProt Gene Name	BTLA
UniProt Entry Name	BTLA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-active-protein/btla/257998>