

Oncostatin M/OSM Active Protein | OSM active protein

Oncostatin M/OSM Protein, Mouse, Recombinant

CATALOG #
AAA258129



HOST
E Coli

PRODUCT SYNONYMS

Oncostatin M/OSM; Oncostatin M/OSM Protein; Mouse; Recombinant; Mouse Oncostatin M/OSM Protein; OncoM Protein; oncostatin M; OSM active protein

PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Ala24-Arg206
Species	Mouse
Activity	1. Measured by its ability to inhibit proliferation of M1 mouse myeloid leukemia cells. The ED50 for this effect is typically 10-50ng/mL. 2. Measured in a cell proliferation assay using NIH3T3 cells. The ED50 for this effect is typically 8-32ng/mL.
Predicted N Terminal	Met
Tag	Native
Protein Construction	A DNA sequence encoding the mouse OSM (P53347)(Ala24-Arg206) was expressed with an initial Met at the N-terminal.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data
Measured in a cell proliferation assay using NIH3T3 cells. The ED50 for this effect is typically 8-32 ng/mL.

ADDITIONAL INFORMATION

Related Product Information for OSM active protein

Background: Oncostatin M (OSM) is a glycoprotein belonging to the interleukin-6 family of cytokines that has functions mainly in cell growth. Oncostatin M (OSM) is considered as a pleiotropic cytokine that signals through cell surface receptors type I and type II both of which share the similarity of containing protein gp130 and takes part in many bio metabolism processes including liver development, hematopoiesis, inflammation, bone formation, and destruction and possibly CNS development. Oncostatin M (OSM) was previously identified by its ability to inhibit the growth of cells from melanoma and other solid tumors. It also has been reported that OSM, like LIF, IL-6, and G-CSF, can inhibit the proliferation of murine M1 myeloid leukemic cells and can induce their differentiation into macrophage-like cells. The human form of OSM is insensitive between pH2 and 11 and resistant to heating for one hour at 56 degrees but is not stable at 90 degrees. The human OSM is produced as a precursor containing 252 amino acids, whose first 25 amino acids function as a secretory signal peptide and which on removal yields the soluble 227 amino acid pro-OSM. Removal of the C-terminal most 31 amino acids produces the fully active 196 residue form.

References

Tanaka M, et al. (2003) Oncostatin M, a multifunctional cytokine. Rev Physiol Biochem Pharmacol. Reviews of Physiology, Biochemistry and Pharmacology. 149: 39-52. Auguste P, et al. (1997) Signaling of type II oncostatin M receptor. J Biol Chem. 272 (25): 15760-4. Zarling JM, et al. (1986). Oncostatin M: a growth regulator produced by differentiated histiocytic lymphoma cells. Proc Natl Acad Sci. 83 (24): 9739-43.

NCBI AND UNIPROT INFORMATION

NCBI GI #	61656173
NCBI GeneID	18413
NCBI Accession #	NP_001013383.1
NCBI GenBank Nucleotide #	NM_001013365.2
UniProt Accession #	P53347; Q3U1Y5; Q5SPX6
Molecular Weight	30,114 Da
NCBI Official Full Name	oncostatin-M
NCBI Official Synonym Full Names	oncostatin M
NCBI Official Symbol	Osm
NCBI Official Synonym Symbols	OncoM
NCBI Protein Information	oncostatin-M
UniProt Protein Name	Oncostatin-M

UniProt Gene Name	Osm
UniProt Synonym Gene Names	OSM
UniProt Entry Name	ONCM_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/oncostatin-m-osm/258129>