

uPAR/PLAUR Active Protein | PLAUR active protein

uPAR/PLAUR Protein, Mouse, Recombinant (His Tag)

CATALOG #
AAA258135


HOST
HEK293 Cells

PRODUCT SYNONYMS

uPAR/PLAUR; uPAR/PLAUR Protein; Mouse; Recombinant (His Tag); Mouse PLAUR/CD87/uPAR Protein (His Tag); Cd87 Protein; u-PAR Protein; uPAR Protein; plasminogen activator; urokinase receptor; PLAUR active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>97% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Thr29 7
Species	Mouse
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Leu 24
Tag	C-His
Protein Construction	A DNA sequence encoding the extracellular domain of mouse PLAUR (NP_035243.1) precursor (Met 1-Thr 297) was expressed, fused with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Activity	Measured by its ability to bind with uPA-His in a functional ELISA.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data
Measured by its ability to bind with uPA-His in a functional ELISA.

ADDITIONAL INFORMATION

Related Product Information for PLAUR active protein	Background: Urokinase plasminogen activator (uPA) and/or its receptor (uPAR) are essential for metastasis, and overexpression of these molecules is strongly correlated with poor prognosis in a variety of malignant tumours. uPAR and uPA levels in both resected tumor tissue and plasma are of independent prognostic significance for patient survival in several types of human cancer. This system has classically been thought to drive tumor progression by mediating directed extracellular proteolysis on the surface of migrating or invading cells, and intervening with this proteolysis by targeting uPAR has been proposed to represent a novel approach for inhibiting tumor progression. uPAR, also known as PLAUR or CD87, has been implicated in the growth, metastasis, and angiogenesis of several solid and hemotologic malignancies. uPAR is a highly glycosylated, 55-60kDa integral membrane protein linked to the plasma membrane by a glycosylphosphatidylinositol (GPI) anchor. It is part of a cell surface system that also consists of the serine protease uPA and several specific inhibitors (plasminogen activator inhibitors 1 and 2). Additionally, the analysis of CD87 (urokinase-type plasminogen activator receptor - uPAR) expression has a potential role in the test or prognostic work-up of several hematological malignancies, particularly acute leukemia and multiple myeloma.
Product Categories/Family for PLAUR active protein	Serine Proteases and Regulators
References	Romer J, et al. (2004) The urokinase receptor as a potential target in cancer therapy. Curr Pharm Des. 10(19): 2359-76. Bn MC, et al. (2004) CD87 (urokinase-type plasminogen activator receptor), function and pathology in hematological disorders: a review. Leukemia. 18(3): 394-400. Pillay V, et al. (2007) The urokinase plasminogen activator receptor as a gene therapy target for cancer. Trends Biotechnol. 25(1): 33-9. Mazar AP. (2008) Urokinase plasminogen activator receptor choreographs multiple ligand interactions: implications for tumor progression and therapy. Clin Cancer Res. 14(18): 5649-55.

NCBI AND UNIPROT INFORMATION

NCBI GI #	7242179
NCBI GeneID	18793
NCBI Accession #	NP_035243.1
NCBI GenBank Nucleotide #	NM_011113.3
UniProt Accession #	P35456; P35457
Molecular Weight	160,000
NCBI Official Full Name	urokinase plasminogen activator surface receptor
NCBI Official Synonym Full Names	plasminogen activator, urokinase receptor
NCBI Official Symbol	Plaur
NCBI Official Synonym Symbols	Cd87; uPAR; u-PAR

NCBI Protein Information	urokinase plasminogen activator surface receptor; urokinase plasminogen activator receptor; urokinase-type plasminogen activator receptor
UniProt Protein Name	Urokinase plasminogen activator surface receptor
UniProt Gene Name	Plaur
UniProt Synonym Gene Names	U-PAR; uPAR
UniProt Entry Name	UPAR_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-active-protein/upar-plaur/258135>