

ICOS active protein

ICOS Protein, Mouse, Recombinant (His Tag)

CATALOG #
AAA258165

HOST
HEK293 Cells



PRODUCT SYNONYMS

ICOS; ICOS Protein; Mouse; Recombinant (His Tag); Mouse ICOS Protein (His Tag); AILIM Protein; CCLP Protein; CRP-1 Protein; H4 Protein; Ly115 Protein; inducible T-cell co-stimulator; ICOS active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>90% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Gln13 5
Species	Mouse
Endotoxin	<1.0EU per ug protein as determined by the LAL method.
Predicted N Terminal	Glu 21
Tag	C-His
Protein Construction	A DNA sequence encoding the mouse ICOS (NP_034643.2)(Met1-Gln135) was expressed with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Activity	Measured by its binding ability in a functional ELISA. Immobilized Rat ICOS ligand hFc at 2ug/mL (100uL/well) can bind Mouse ICOS His. The EC50 of Mouse ICOS His is 150-800ng/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.

IMAGES



SDS-PAGE

Application Data
Measured by its binding ability in a functional ELISA. Immobilized Rat ICOS ligand hFc at 2 ug/mL (100 uL/well) can bind Mouse ICOS His, the EC50 of Mouse ICOS His is 150-800 ng/mL.

ADDITIONAL INFORMATION

Related Product Information for ICOS active protein	Background: Inducible costimulator (ICOS), also called AILIM (Activation-Inducible Lymphocyte Immunomediatory Molecule) is a cell-surface receptor and belongs to the CD28 family of immune costimulatory receptors consisting of CD28, CTLA-4, and PD-1. The interaction of B7-H2/ICOS plays a critical role in Th cell differentiation, T-B cell interactions which are essential for the germinal center formation, and humoral immune responses, and as well as the production of cytokine IL-4. Also, ICOS is more potent in the induction of IL-10 production, a cytokine important for the suppressive function of T regulatory cells. The B7-1/B7-2--CD28/CTLA-4 and ICOS-B7RP-1 pathway provide key second signals that can regulate the activation, inhibition, and fine-tuning of T-lymphocyte responses. ICOS stimulates both Th1 and Th2 cytokine production but may have a preferential role in Th2 cell development. Moreover, The B7-1/B7-2-CD28/CTLA-4 and ICOS-B7RP-1 pathway has been suggested as being involved in the development of airway inflammation and airway hyperresponsiveness.
Product Categories/Family for ICOS active protein	Cancer Drug Targets
References	Coyle AJ, et al. (2004) The role of ICOS and other costimulatory molecules in allergy and asthma. Springer Semin Immunopathol. 25(3-4): 349-59.Chen YQ, et al. (2006) CD28/CTLA-4--CD80/CD86 and ICOS--B7RP-1 costimulatory pathway in bronchial asthma. Allergy. 61(1): 15-26.van Berkel ME, et al. (2006) CD28 and ICOS: similar or separate costimulators of T cells Immunol Lett. 105(2): 115-22.

NCBI AND UNIPROT INFORMATION

NCBI GI #	224809335
NCBI GeneID	54167
NCBI Accession #	NP_059508.2
NCBI GenBank Nucleotide #	NM_017480.2
UniProt Accession #	Q9WVS0; Q9JL17
Molecular Weight	22,709 Da
NCBI Official Full Name	inducible T-cell costimulator
NCBI Official Synonym Full Names	inducible T cell co-stimulator
NCBI Official Symbol	Icos
NCBI Official Synonym Symbols	H4; CCLP; AILIM; CRP-1; Ly115
NCBI Protein Information	inducible T-cell costimulator
UniProt Protein Name	Inducible T-cell costimulator

UniProt Gene Name	Icos
UniProt Synonym Gene Names	Ailim; CCLP; CRP-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-active-protein/icos/258165>