

Epiregulin Active Protein | EREG active protein

Epiregulin Protein, Mouse, Recombinant (hFc Tag)

CATALOG #
AAA258183


HOST
HEK293 Cells

PRODUCT SYNONYMS

Epiregulin; Epiregulin Protein; Mouse; Recombinant (hFc Tag); Mouse Epiregulin/EREG Protein (Fc Tag); EPR Protein; epiregulin; EREG active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Val56-Leu1 01
Species	Mouse
Activity	Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is typically 0.5-2.5ug/mL.
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Glu
Tag	N-human IgG1-Fc
Protein Construction	A DNA sequence encoding the mature form of mouse EREG (Q61521)(Val 56-Leu 101) was fused with the Fc region of human IgG1 at the N-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data
Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is typically 0.5-2.5 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for EREG active protein	Background: Epiregulin (EREG) is a member of the epidermal growth factor family. Epiregulin (EREG) can function as a ligand of EGFR (epidermal growth factor receptor), as well as a ligand of most members of the ERBB (v-erb-b2 oncogene homolog) family of tyrosine-kinase receptors. Epiregulin (EREG) exhibit bifunctional regulatory properties: it inhibit the growth of several epithelial tumor cells and stimulated the growth of fibroblasts and various other types of cells. Epiregulin (EREG) bound to the EGF receptors of epidermoid carcinoma A431 cells much more weakly than did EGF, but was nevertheless much more potent than EGF as a mitogen for rat primary hepatocytes and Balb/c 3T3 A31 fibroblasts. These findings suggest that epiregulin (EREG) plays important roles in regulating the growth of epithelial cells and fibroblasts by binding to receptors for EGF-related ligands. Epiregulin (EREG) is the broadest specificity EGF-like ligand so far characterized: not only does it stimulate homodimers of both ErbB-1 and ErbB-4, it also activates all possible heterodimeric ErbB complexes.
References	Shelly M, et al. (1998) Epiregulin is a potent pan-ErbB ligand that preferentially activates heterodimeric receptor complexes. J Biol Chem. 1998 Apr 24;273(17):10496-505. Shirakata Y, et al. (2000) Epiregulin, a novel member of the epidermal growth factor family, is an autocrine growth factor in normal human keratinocytes. J Biol Chem. 275(8): 5748-53. Zhu Z, et al. (2000) Epiregulin is Up-regulated in pancreatic cancer and stimulates pancreatic cancer cell growth. Biochem Biophys Res Commun. 273(3): 1019-24.

NCBI AND UNIPROT INFORMATION

NCBI GI #	6679683
NCBI GeneID	13874
NCBI Accession #	NP_031976.1
NCBI GenBank Nucleotide #	NM_007950.2
UniProt Accession #	Q61521; Q7M0A8
Molecular Weight	18,474 Da
NCBI Official Full Name	proepiregulin preproprotein
NCBI Official Synonym Full Names	epiregulin
NCBI Official Symbol	Ereg
NCBI Official Synonym Symbols	EPR
NCBI Protein Information	proepiregulin
UniProt Protein Name	Proepiregulin
UniProt Gene Name	Ereg

UniProt Synonym Gene Names	EPR
UniProt Entry Name	EREG_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-ereg-active-protein/epiregulin/258183>