

PD-L2 Active Protein | PDCD1LG2 active protein

PD-L2 Protein, Mouse, Recombinant (hFc Tag)

CATALOG #
AAA258204


HOST
HEK293 Cells

PRODUCT SYNONYMS

PD-L2; PD-L2 Protein; Mouse; Recombinant (hFc Tag); Mouse B7-DC/PD-L2/CD273 Protein (Fc Tag); B7-DC Protein; Btdc Protein; F730015O22Rik Protein; programmed cell death 1 ligand 2; PDCD1LG2 active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-Arg21 9
Species	Mouse
Endotoxin	<1.0EU per ug of the protein as determined by the LAL method
Predicted N Terminal	Leu 20
Tag	C-human IgG1-Fc
Protein Construction	A DNA sequence encoding the extracellular domain of mouse PDCD1LG2 (Q9WUL5)(Met 1-Arg 219) was fused with the Fc region of human IgG1 at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Activity	1. Measured by its binding ability in a functional ELISA. 2. Immobilized human PD1-His at 10ug/mL (100uL/well) can bind mouse PD-L2-Fc. The EC50 of mouse PD-L2-Fc is 0.95ug/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.



SDS-PAGE

Application Data
Measured by its binding ability in a functional ELISA. Immobilized human PD1-His at 10 ug/mL (100 ul/well) can bind mouse PD-L2-Fc, The EC50 of mouse PD-L2-Fc is 0.95 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for PDCD1LG2 active protein	Background: Programmed death ligand 2 (PD-L2), also referred to as B7-DC and CD273, is a member of the B7 family of proteins including B7-1, B7-2, B7-H2, B7-H1 (PD-L1), and B7-H3. PD-L2 is a type I membrane protein and structurally consists of an extracellular region containing one V-like and one C-like Ig domain, a transmembrane region, and a short cytoplasmic domain. PD-L2 is expressed on antigen presenting cells, placental endothelium and medullary thymic epithelial cells, and can be induced by LPS in B cells, INF-? in monocytes, or LPS plus IFN-? in dendritic cells. The CD28 and B7 protein families are critical regulators of immune responses. PD-L2 and PD-L1 are two ligands for PD-1, member of the CD28/CTLA4 family expressed on activated lymphoid cells, and thus provide signals for regulating T cell activation and immune tolerance. The interaction of B7-DC/PD-1 exhibited a 2-6-fold higher affinity compared with the interaction of B7-H1/PD-1.
Product Categories/Family for PDCD1LG2 active protein	Cancer Drug Targets
References	Latchman Y, et al. (2001) PD-L2 is a second ligand for PD-1 and inhibits T cell activation. Nat Immunol. 2: 261-8.Carreno BM, et al. (2005) Therapeutic opportunities in the B7/CD28 family of ligands and receptors. Curr Opin Pharmacol. 5(4): 424-30.Radhakrishnan S, et al. (2007) B7-DC/PD-L2 cross-linking induces NF-kappaB-dependent protection of dendritic cells from cell death. J Immunol. 178(3): 1426-32.

NCBI AND UNIPROT INFORMATION

NCBI GI #	10946740
NCBI GeneID	58205
NCBI Accession #	NP_067371.1
NCBI GenBank Nucleotide #	NM_021396.2
UniProt Accession #	Q9WUL5
Molecular Weight	27,820 Da
NCBI Official Full Name	programmed cell death 1 ligand 2
NCBI Official Synonym Full Names	programmed cell death 1 ligand 2
NCBI Official Symbol	Pdcd1lg2
NCBI Official Synonym Symbols	Btdc; B7-DC; PD-L2; F730015O22Rik
NCBI Protein Information	programmed cell death 1 ligand 2; PD-1 ligand 2; PDCD1 ligand 2; butyrophilin B7-DC; butyrophilin-like protein; programmed death ligand 2
UniProt Protein Name	Programmed cell death 1 ligand 2

UniProt Gene Name	Pdcd1lg2
UniProt Synonym Gene Names	B7dc; Btdc; Cd273; Pdl2; PD-1 ligand 2; PD-L2; PDCD1 ligand 2; Programmed death ligand 2; B7-DC
UniProt Entry Name	PD1L2_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-pdcd1lg2-active-protein/pd-l2/258204>