

DKK1 active protein

DKK1 Protein, Mouse, Recombinant (His Tag)

CATALOG #
AAA258244

HOST
HEK293 Cells


PRODUCT SYNONYMS

DKK1; DKK1 Protein; Mouse; Recombinant (His Tag); Mouse DKK-1 Protein (His Tag); mdck-1 Protein; dickkopf WNT signaling pathway inhibitor 1; DKK1 active protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the hard copy of CoA.
Sequence	Met1-His27 2
Species	Mouse
Activity	Measured by its ability to inhibit recombinant rmWnt3a induced alkaline phosphatase production by C3H10T 1/2 cells. The ED50 for this effect is 0.05-0.3ug/mL.
Endotoxin	<1.0EU per ug protein as determined by the LAL method.
Predicted N Terminal	Thr 32
Tag	C-His
Protein Construction	A DNA sequence encoding the mouse DKK1 (NP_034181.2)(Met1-His272) was expressed with a polyhistidine tag at the C-terminus.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice.

IMAGES



SDS-PAGE

Application Data
Measured by its ability to inhibit recombinant rmWnt3a induced alkaline phosphatase production by C3H10T 1/2 cells. The ED50 for this effect is 0.05-0.3 ug/mL.

ADDITIONAL INFORMATION

Related Product Information for DKK1 active protein	Background: Dickkopf (DKK) family proteins, consisting of DKK-1, DKK-2, DKK-3 and DKK-4, function as secreted Wnt antagonists by inhibiting Wnt coreceptors LRP5/6. DKK-1, DKK-2, and DKK-4 also bind cell surface Kremen-1 or Kremen-2 and promote the internalization of LRP5/6. Dickkopf related protein 1 (DKK-1) was initially identified as an inducer of head formation in Xenopus embryos. DKK-1 protein modulates Wnt signaling pathway during embryonic development. Increased levels of DKK-1 are found in the majority of lung cancers, esophageal squamous cell carcinomas, and hormone-resistant breast cancers, while DKK-1 expression is decreased in malignant melanoma and colorectal cancers.
Product Categories/Family for DKK1 active protein	Canonical Wnt Pathway
References	Horwitz EM. (2004) Dkk-1-mediated expansion of adult stem cells. Trends Biotechnol. 22(8): 386-8.Jiang T, et al. (2009) Clinical significance of serum DKK-1 in patients with gynecological cancer. Int J Gynecol Cancer. 19(7): 1177-81.

NCBI AND UNIPROT INFORMATION

NCBI GI #	31542557
NCBI GeneID	13380
NCBI Accession #	NP_034181.2
NCBI GenBank Nucleotide #	NM_010051.3
UniProt Accession #	O54908; Q80UL5
Molecular Weight	29,298 Da
NCBI Official Full Name	dickkopf-related protein 1
NCBI Official Synonym Full Names	dickkopf WNT signaling pathway inhibitor 1
NCBI Official Symbol	Dkk1
NCBI Official Synonym Symbols	mdkk-1
NCBI Protein Information	dickkopf-related protein 1
UniProt Protein Name	Dickkopf-related protein 1
UniProt Gene Name	Dkk1
UniProt Synonym Gene Names	Dickkopf-1; Dkk-1; mDkk-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/hek293-cells-active-protein/dkk1/258244>