

Rabbit anti-Human FAM129B Polyclonal Antibody | anti-FAM129B antibody

Anti-FAM129B Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258506	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry

PRODUCT SYNONYMS

FAM129B; Polyclonal Antibody; Anti-FAM129B Antibody; Rabbit Polyclonal; FAM129B Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-bA356B19.6 Antibody; Anti-C9orf88 Antibody; Anti-MEG-3 Antibody; Anti-MINERVA Antibody; Anti-OC58 Antibody; OC58; MEG-3; C9orf88; MINERVA; bA356B19.6; anti-FAM129B antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FAM129B
Form/Format	PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FAM129B antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	E. coli-derived Human FAM129B fragment
Preparation	Produced in rabbits immunized with E. coli-derived Human FAM129B fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescence staining of FAM129B in U251MG cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human FAM129B polyclonal antibody (dilution ratio 1:100) at 4° overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Cell membrane.



IHC (Immunohistochemistry)

Immunohistochemical staining of human FAM129B in human stomach with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunohistochemical staining of human FAM129B in human placenta with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

References	Lee JH, et al. (2016) Fam129b activates ras and promotes aerobic glycolysis. Cell Cycle 15 (11): 1391-1392.Ji H, et al. (2016) Egfr phosphorylates fam129b to promote ras activation. Proc Natl Acad Sci U S A 113 (3): 644-649.
------------	--

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fam129b-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/258506>