

COVID 19 (AY.3) Spike S1+S2 trimer Coronavirus Recombinant Protein | COVID-19 recombinant protein

SARS-CoV-2 (AY.3) Spike S1+S2 trimer Protein (ECD, His tag)



CATALOG # AAA258553	HOST HEK293 Cells	REACTIVITY SARS-CoV-2
-------------------------------	-----------------------------	---------------------------------

PRODUCT SYNONYMS

COVID 19 (AY.3) Spike S1+S2 trimer Coronavirus; SARS-CoV-2 (AY.3) Spike S1+S2 trimer Protein (ECD; His tag); 2019 Novel Coronavirus; Coronavirus; CoV; COVID-19 virus; HCoV-2; Human Coronavirus 2019; SARS2; SARS-CoV-2; Severe acute respiratory syndrome coronavirus 2; (AY.3) Spike S1+S2 trimer; COVID-19 recombinant protein


PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Reactivity	SARS-CoV-2
Purity/Purification	>95% as determined by SDS-PAGE. >90% as determined by SEC-HPLC.
Form/Format	Lyophilized from sterile 25mM Sodium Citrate, 200mM NaCl, 0.02% tween80, pH6.0. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Sequence	Met1-Gln12 08
Endotoxin	<1.0EU per ug protein as determined by the LAL method.
Tag	C-His
Predicted N Terminal	Val 16
Protein Construction	A DNA sequence encoding the SARS-CoV-2 Spike S1+S2 (YP_009724390.1, with mutations T19R, E156G, DEL157-158, L452R, T478K, D614G, P681R, D950N, F817P, A892P, A899P, A942P, K986P, V987P and furin cleavage site mutants) expressed with the bacteriophage T4 fibrin and a polyhistidine tag at the C-terminus. The mutations were identified in the SARS-CoV-2 variant (known as variant AY.3, AY.3.1) which emerged in the India.
Activity	Immobilized human ACE2 protein (mFc tag) at 2 ug/mL (100 uL/well) can bind SARS-CoV-2 (AY.3) Spike S1+S2 trimer Protein (ECD, His tag), the EC50 is 6-32 ng/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise.



Application Data

Application Data
Immobilized human ACE2 protein (mFc tag)(Cat:10108-H05H) at 2 ug/mL (100 uL/well) can bind SARS-CoV-2 (AY.3) Spike S1+S2 trimer Protein (ECD, His tag)(Cat:AAA258553), the EC50 is 6-32 ng/mL.



Application Data

ADDITIONAL INFORMATION

Related Product Information for COVID-19 recombinant protein

The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.

References

Shen S, et al. (2007) Expression, glycosylation, and modification of the spike (S) glycoprotein of SARS CoV. Methods Mol Biol. 379: 127-35. Du L, et al. (2009) The spike protein of SARS-CoV--a target for vaccine and therapeutic development. Nat Rev Microbiol. 7 (3): 226-36. Xiao X, et al. (2004) The SARS-CoV S glycoprotein. Cell Mol Life Sci. 61 (19-20): 2428-30.

NCBI AND UNIPROT INFORMATION

Molecular Weight	51.5 kDa
------------------	----------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/covid-19-ay-3-spike-s1-s2-trimer-coronavirus-recombinant-protein-hek293-cells/258553>