

Rabbit anti-Human EIF4A3 Polyclonal Antibody | anti-EIF4A3 antibody

EIF4A3 Antibody, Rabbit PAb, Antigen Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258647	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

EIF4A3; Polyclonal Antibody; EIF4A3 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-DDX48 Antibody; Anti-eIF4AIII Antibody; Anti-MUK34 Antibody; Anti-NMP265 Antibody; Anti-NUK34 Antibody; Anti-RCPS Antibody; anti-EIF4A3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Human EIF4A3
Applicable Applications for anti-EIF4A3 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Preparation	Produced in rabbits immunized with E Coli-derived Human EIF4A3 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IP (Immunoprecipitation)

EIF4A3 was immunoprecipitated using;; Lane A:0.5 mg HeLa Whole Cell Lysate; Lane B:0.5 mg MCF7 Whole Cell Lysate; Lane C:0.5 mg Raji Whole Cell Lysate; Lane D:0.5 mg HepG2 Whole Cell Lysate; 4 uL anti-EIF4A3 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G.; Primary antibody;; Anti-EIF4A3 rabbit polyclonal antibody,at 1:100 dilution; Secondary antibody;; Clean-Blot IP Detection Reagent (HRP) at 1:1000dilution; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size: 45 kDa; Observed band size :45 kDa




IHC (Immunohistochemistry)

Immunochemical staining of human EIF4A3 in human kidney with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human EIF4A3 in human liver with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-EIF4A3 rabbit polyclonal antibody at 1:500 dilution; Lane A: HeLa Whole Cell Lysate; Lane B: MCF7 Whole Cell Lysate; Lane C: Raji Whole Cell Lysate; Lane D: HepG2 Whole Cell Lysate; Lysates/proteins at 30 ug per lane.; Secondary; Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size:45 kDa; Observed band size:45 kDa

ADDITIONAL INFORMATION

Product Categories/Family for anti-EIF4A3 antibody	Neuroscience
--	--------------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/eif4a3-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/258647>