

Rabbit anti-Human HIGD1A Polyclonal Antibody | anti-HIGD1A antibody

HIGD1A Antibody, Rabbit PAb, Antigen Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258659	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

HIGD1A; Polyclonal Antibody; HIGD1A Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-HIG1 Antibody; Anti-RCF1a Antibody; anti-HIGD1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Human HIGD1A
Applicable Applications for anti-HIGD1A antibody	IP (Immunoprecipitation), WB (Western Blot)
Preparation	Produced in rabbits immunized with E Coli-derived Human HIGD1A fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

HIGD1A was immunoprecipitated using;; Lane A:0.5 mg HepG2 Whole Cell Lysate; 4 uL anti-HIGD1A rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G.; Primary antibody;; Anti-HIGD1A rabbit polyclonal antibody,at 1:100 dilution; Secondary antibody;; Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size: 10 kDa; Observed band size :10 kDa



WB (Western Blot)

Anti-HIGD1A rabbit polyclonal antibody at 1:500 dilution; Lane A: A431 Whole Cell Lysate; Lane B: Hela Whole Cell Lysate; Lane C: HepG2 Whole Cell Lysate; Lane D: 293 Whole Cell Lysate; Lysates/proteins at 30 ug per lane.; Secondary; Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.; Developed using the ECL technique.; Performed under reducing conditions.; Predicted band size:12 kDa; Observed band size:12 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.