

Rabbit anti-Human CAV2/Caveolin-2 Polyclonal Antibody | anti-CAV2 antibody

Anti-CAV2/Caveolin-2 Antibody, Rabbit Polyclonal

CATALOG # AAA258724	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CAV2/Caveolin-2; Polyclonal Antibody; Anti-CAV2/Caveolin-2 Antibody; Rabbit Polyclonal; caveolin 2; Anti-CAV Antibody; anti-CAV2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CAV2/Caveolin-2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CAV2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human CAV2/Caveolin-2
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human CAV2/Caveolin-2, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

CAV2 was immunoprecipitated using: Lane A: 0.5 mg A431 Whole Cell Lysate Lane B: 0.5 mg HeLa Whole Cell Lysate 4 uL anti-CAV2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CAV2 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 18 kDa Observed band size: 18 kDa



WB (Western Blot)

Anti-CAV2/Caveolin-2 rabbit polyclonal antibody at 1:500 dilution Lane A: Huvec Whole Cell Lysate Lane B: A431 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 18 kDa Observed band size: 18 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cav2-caveolin-2-antibody-immunoprecipitation-western-blot-elisa/258724>