

Rabbit anti-Human PRMT6 Polyclonal Antibody | anti-PRMT6 antibody

Anti-PRMT6 Antibody, Rabbit Polyclonal

CATALOG # AAA258745	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

PRMT6; Polyclonal Antibody; Anti-PRMT6 Antibody; Rabbit Polyclonal; protein arginine methyltransferase 6; Anti-HRMT1L6 Antibody; anti-PRMT6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human PRMT6
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-PRMT6 antibody	IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human PRMT6
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human PRMT6, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human PRMT6 in human carcinoma of sigmoid with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human PRMT6 in human breast carcinoma with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.

ADDITIONAL INFORMATION

Related Product Information for anti-PRMT6 antibody	Protein arginine N-methyltransferase 6, also known as Histone-arginine N-methyltransferase PRMT6, PRMT6, and HRMT1L6, is a member of the protein arginine N-methyltransferase family and PRMT6 subfamily. PRMT6 is highly expressed in kidney and testes. PRMT6 is known to catalyze the generation of asymmetric dimethylarginine in polypeptides. It has been implicated in human immunodeficiency virus pathogenesis, DNA repair, and transcriptional regulation. PRMT6 is known to methylate histone H3 Arg-2 (H3R2), and this negatively regulates the lysine methylation of H3K4 resulting in gene repression. PRMT6 plays a key role in coupling process by functioning as a transcriptional coactivator that can regulate alternative splicing. PRMT6 coactivates the progesterone, glucocorticoid and oestrogen receptors in luciferase reporter assays in a hormone-dependent manner. Small interfering RNA (siRNA) oligonucleotide duplex knockdown of PRMT6 disrupts oestrogen-stimulated transcription of endogenous GREB1 and progesterone receptor in MCF-7 breast cancer cells. Neutralizing the activity of PRMT6 could inhibit tumor progression and may be of cancer therapeutic significance.
References	Hyllus D, et al., 2007, Genes Dev. 21(24): 3369-80. Lakowski, TM. et al., 2008, J Biol Chem. 283 (15): 10015-25. Michaud-Levesque, J. et al., 2009, J Biol Chem. 284 (32): 21338-46. Harrison, MJ. et al., 2010, Nucleic acids Res. Jan 4.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/prmt6-antibody-immunohistochemistry-elisa/258745>