

Rabbit anti-Human CPSF6 Polyclonal Antibody | anti-CPSF6 antibody

Anti-CPSF6 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258790	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CPSF6; Polyclonal Antibody; Anti-CPSF6 Antibody; Rabbit Polyclonal; cleavage and polyadenylation specific factor 6; Anti-CFIM Antibody; Anti-CFIM68 Antibody; Anti-HPBRII-4 Antibody; Anti-HPBRII-7 Antibody; anti-CPSF6 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CPSF6
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-CPSF6 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the C-terminus of the Human CPSF6
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the C-terminus of the Human CPSF6, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES




IP (Immunoprecipitation)

CPSF6 was immunoprecipitated using: Lane A: 0.5 mg Jurkat Whole Cell Lysate; Lane B: 0.5 mg K562 Whole Cell Lysate; 4 uL anti-CPSF6 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CPSF6 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 59 kDa. Observed band size: 72 kDa.




IHC (Immunohistochemistry)

Immunohistochemical staining of human CPSF6 in human carcinoma of sigmoid with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human CPSF6 in human breast carcinoma with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-CPSF6 rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lane B: Jurkat Whole Cell Lysate Lane C: NIH-3T3 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 59 kDa Observed band size: 70 kDa (We are unsure as to the identity of these extra bands).

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cpsf6-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/258790>