

Rabbit anti-Human IRF2/IRF-2 Polyclonal Antibody | anti-IRF2/IRF-2 antibody

Anti-IRF2/IRF-2 Antibody, Rabbit Polyclonal


CATALOG # AAA258805	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

IRF2/IRF-2; Polyclonal Antibody; Anti-IRF2/IRF-2 Antibody; Rabbit Polyclonal; interferon regulatory factor 2; Anti-IRF-2 Antibody; anti-IRF2/IRF-2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human IRF2/IRF-2
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-IRF2/IRF-2 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	A synthetic peptide corresponding to the of the Human IRF2/IRF-2
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the of the Human IRF2/IRF-2, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.



IP (Immunoprecipitation)

IRF2 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate Lane B: 0.5 mg 293T Whole Cell Lysate 4 uL anti-IRF2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-IRF2 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Clean-Blot IP Detection Reagent (HRP) at 1:1000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 39 kDa Observed band size :45 kDa



WB (Western Blot)

Anti-IRF2 rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lane B: 293T Whole Cell Lysate Lane C: HepG2 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 39 kDa Observed band size: 45 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.