

Rabbit anti-Human LysRS/Lysyl tRNA synthetase Polyclonal Antibody | anti-LysRS antibody

Anti-LysRS/Lysyl tRNA synthetase Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258810	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry


PRODUCT SYNONYMS

LysRS/Lysyl tRNA synthetase; Polyclonal Antibody; Anti-LysRS/Lysyl tRNA synthetase Antibody; Rabbit Polyclonal; lysyl-tRNA synthetase; Anti-CMTRIB Antibody; Anti-DFNB89 Antibody; Anti-KARS1 Antibody; Anti-KARS2 Antibody; Anti-KRS Antibody; anti-LysRS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human LysRS/Lysyl tRNA synthetase
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-LysRS antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry)
Immunogen	A synthetic peptide corresponding to the N-terminus of the Human LysRS/Lysyl tRNA synthetase
Preparation	Produced in rabbits immunized with a synthetic peptide corresponding to the N-terminus of the Human LysRS/Lysyl tRNA synthetase, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of HPV-SMCCA5875+HPV-SMCCA5296 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human HPV-SMCCA5875+HPV-SMCCA5296 polyclonal antibody (dilution ratio 1:100) at 4? overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunohistochemical staining of human KARS in human breast carcinoma with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human KARS in human colon with rabbit polyclonal antibody at 1:500 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/lysrs-lysyl-trna-synthetase-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-elisa/258810>