

Rabbit anti-Human SF20 Polyclonal Antibody | anti-SF20 antibody

Anti-SF20 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258910	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

SF20; Polyclonal Antibody; Anti-SF20 Antibody; Rabbit Polyclonal; myeloid-derived growth factor; Anti-C19orf10 Antibody; Anti-EUROIMAGE1875335 Antibody; Anti-IL25 Antibody; Anti-IL27 Antibody; Anti-IL27w Antibody; Anti-R33729_1 Antibody; anti-SF20 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human SF20
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; 0.2um filtered solution in PBS
Applicable Applications for anti-SF20 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant Human SF20 Protein
Preparation	Produced in rabbits immunized with purified, recombinant Human SF20 (rh SF20). SF20 specific IgG was purified by Human SF20 affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Preservative-Free. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

Lane A: 0.5 mg 293T Whole Cell Lysate
4 uL anti-C19ORF10 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G.
Primary antibody: Anti-C19ORF10 rabbit polyclonal antibody, at 1:100 dilution
Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution
Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 15 kDa
Observed band size: 15 kDa



IHC (Immunohistochemistry)

Immunohistochemical staining of human SF20 in human liver with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human SF20 in human pancreas with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-SF20 rabbit polyclonal antibody at 1:500 dilution
Lane A: 293T Whole Cell Lysate
Lane B: MCF7 Whole Cell Lysate
Lysates/proteins at 30 ug per lane.
Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.
Developed using the ECL technique.
Performed under reducing conditions.
Predicted band size: 19 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/sf20-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/258910>