

Rabbit anti-Human CSNK1E/CK1 epsilon Polyclonal Antibody | anti-CSNK1E/CK1epsilon antibody

Anti-CSNK1E/CK1 epsilon Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258934	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

CSNK1E/CK1 epsilon; Polyclonal Antibody; Anti-CSNK1E/CK1 epsilon Antibody; Rabbit Polyclonal; casein kinase 1; epsilon; Anti-CK1epsilon Antibody; Anti-HCK1E Antibody; anti-CSNK1E/CK1epsilon antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CSNK1E/CK1 epsilon
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CSNK1E/CK1epsilon antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived Human CSNK1E fragment
Preparation	Produced in rabbits immunized with E coli-derived Human CSNK1E fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of CSNK1E in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human CSNK1E polyclonal antibody (dilution ratio 1:200) at 4° overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus and Cytoplasm.



WB (Western Blot)

Anti-CSNK1E rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 47 kDa Observed band size: 47 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.