

Rabbit anti-Human Fub1 Polyclonal Antibody | anti-Fub1 antibody

Anti-Fub1 Antibody, Rabbit Polyclonal


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA258982	Rabbit	Human	Western Blot, Immunofluorescence, Immunocytochemistry

PRODUCT SYNONYMS

Fub1; Polyclonal Antibody; Anti-Fub1 Antibody; Rabbit Polyclonal; Finkel-Biskis-Reilly murine sarcoma virus (FBR-MuSV) ubiquitously expressed; Anti-asr1 Antibody; Anti-FAU1 Antibody; Anti-Fubi Antibody; Anti-MNSFbeta Antibody; Anti-RPS30 Antibody; Anti-S30 Antibody; anti-Fub1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human Fub1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-Fub1 antibody	WB (Western Blot), IF (Immunofluorescence), ICC (Immunocytochemistry)
Immunogen	E coli-derived Human Fub1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human Fub1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.



IF (Immunofluorescence)

Immunofluorescence staining of Fub1 in PC3 cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human Fub1 polyclonal antibody (dilution ratio 1:100) at 4° overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Nucleus and Cytoplasm.



WB (Western Blot)

Anti-FAU rabbit monoclonal antibody at 1:500 dilution Lane A: HepG2 Whole Cell Lysate Lane B: HeLa Whole Cell Lysate Lane C: SW480 Whole Cell Lysate Lane D: 293 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 14 kDa Observed band size: 14 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.