

Rabbit anti-Human BTF3 Polyclonal Antibody | anti-BTF3 antibody

Anti-BTF3 Antibody, Rabbit Polyclonal

CATALOG # AAA258998	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	----------------------------	--

PRODUCT SYNONYMS

BTF3; Polyclonal Antibody; Anti-BTF3 Antibody; Rabbit Polyclonal; basic transcription factor 3; Anti-BETA-NAC Antibody; Anti-BTF3a Antibody; Anti-BTF3b Antibody; Anti-NACB Antibody; anti-BTF3 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human BTF3
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-BTF3 antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	E coli-derived Human BTF3 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human BTF3 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

BTF3 was immunoprecipitated using: Lane A: 0.5 mg MCF7 Whole Cell Lysate 4 uL anti-BTF3 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-BTF3 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 22 kDa. Observed band size: 22 kDa.



WB (Western Blot)

Anti-BTF3 rabbit monoclonal antibody at 1:500 dilution. Lane A: MCF7 Whole Cell Lysate. Lane B: HeLa Whole Cell Lysate. Lane C: Jurkat Whole Cell Lysate. Lane D: K562 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 22 kDa. Observed band size: 18 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.