

Rabbit anti-Human EHHADH Polyclonal Antibody | anti-EHHADH antibody

Anti-EHHADH Antibody, Rabbit Polyclonal

CATALOG # AAA259000	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

EHHADH; Polyclonal Antibody; Anti-EHHADH Antibody; Rabbit Polyclonal; enoyl-CoA; hydratase/3-hydroxyacyl CoA dehydrogenase; Anti-ECHD Antibody; Anti-FRTS3 Antibody; Anti-L-PBE Antibody; Anti-LBFP Antibody; Anti-LBP Antibody; Anti-PBFE Antibody; anti-EHHADH antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human EHHADH
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-EHHADH antibody	IHC (Immunohistochemistry)
Immunogen	E coli-derived Human EHHADH fragment
Preparation	Produced in rabbits immunized with E coli-derived Human EHHADH fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human EHHADH in human liver with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human EHHADH in human kidney with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IP (Immunoprecipitation)

EHHADH was immunoprecipitated using: Lane A: 0.5 mg U-251 MG Whole Cell Lysate 4 uL anti-EHHADH rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-EHHADH rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 79 kDa. Observed band size: 90 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ehhadh-antibody-immunohistochemistry-elisa/259000>