

Rabbit anti-Human FAM3C Polyclonal Antibody | anti-FAM3C antibody

Anti-FAM3C Antibody, Rabbit Polyclonal

CATALOG # AAA259020	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---

PRODUCT SYNONYMS

FAM3C; Polyclonal Antibody; Anti-FAM3C Antibody; Rabbit Polyclonal; family with sequence similarity 3; member C; Anti-GS3786 Antibody; Anti-ILEI Antibody; anti-FAM3C antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human FAM3C
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-FAM3C antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human FAM3C fragment
Preparation	Produced in rabbits immunized with E coli-derived Human FAM3C fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IHC (Immunohistochemistry)

Immunohistochemical staining of human FAM3C in human brain with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunohistochemical staining of human FAM3C in human small intestine with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-FAM3C rabbit polyclonal antibody at 1:500 dilution. Lane A: A431 Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 25 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-FAM3C antibody	FAM3C, also known as ILEI, is a member of theFAM3 family. It can be detected in resent in most secretory epithelia. FAM3C may be involved in retinal laminar formation. It promotes epithelial to mesenchymal transition. A change in expression of FAM3C has been noted in pancreatic cancer-derived cells. ILEI is overexpressed and/or altered in intracellular localization in multiple human tumors, an event strongly correlated to invasion/EMT, metastasis formation, and survival in human colon and breast cancer.
References	Zhu Y. et al., 2002, Genomics. 80 (2): 144-50.Strausberg RL. et al., 2003, Proc Natl Acad Sci. 99 (26): 16899-903. Quéméner-Redon S. et al., 2003, Nature. 424 (6945): 157-64.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/fam3c-antibody-immunohistochemistry-western-blot-elisa/259020>