

Rabbit anti-Human MAP4 Polyclonal Antibody | anti-MAP4 antibody

Anti-MAP4 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259032	Rabbit	Human	Immunofluorescence, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MAP4; Polyclonal Antibody; Anti-MAP4 Antibody; Rabbit Polyclonal; anti-MAP4 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human MAP4
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-MAP4 antibody	IF (Immunofluorescence), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human MAP4 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human MAP4 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES





IF (Immunofluorescence)

Immunofluorescence staining of MAP4 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human MAP4 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Cytoplasm.



IHC (Immunohistochemistry)

Immunochemical staining of human MAP4 in human kidney with rabbit polyclonal antibody at 1:5000 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human MAP4 in human epididymis with rabbit polyclonal antibody at 1:5000 dilution, formalin-fixed paraffin embedded sections.

WB (Western Blot)

Anti-MAP4 rabbit polyclonal antibody at 1:500 dilution Lane A: RT4 Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 121 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-MAP4 antibody

microtubule-associated protein 4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/map4-antibody-immunofluorescence-immunocytochemistry-immunohistochemistry-western-blot-elisa/259032>