

Rabbit anti-Human MLLT1 Polyclonal Antibody | anti-MLLT1 antibody

Anti-MLLT1 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259052	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry, Western Blot


PRODUCT SYNONYMS

MLLT1; Polyclonal Antibody; Anti-MLLT1 Antibody; Rabbit Polyclonal; Anti-ENL Antibody; Anti-LTG19 Antibody; Anti-YEATS1 Antibody; anti-MLLT1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human MLLT1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-MLLT1 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry), WB (Western Blot)
Immunogen	E coli-derived Human MLLT1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human MLLT1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescence staining of MLLT1 in U251MG cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human MLLT1 polyclonal antibody (dilution ratio 1:200) at 4? overnight. Then cells were stained with the Alexa Fluor488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus.



WB (Western Blot)

Anti-MLLT1 rabbit polyclonal antibody at 1:500 dilution
Lane A: MCF7 Whole Cell Lysate
Lane B: HeLa Whole Cell Lysate
Lane C: Jurkat Whole Cell Lysate
Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 62 kDa Observed band size: 62 kDa (We are unsure as to the identity of these extra bands).

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mlt1-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-western-blot-elisa/259052>