

Rabbit anti-Human TIMM10 Polyclonal Antibody | anti-TIMM10 antibody

Anti-TIMM10 Antibody, Rabbit Polyclonal

CATALOG # AAA259079	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunoprecipitation, Immunohistochemistry
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TIMM10; Polyclonal Antibody; Anti-TIMM10 Antibody; Rabbit Polyclonal; Anti-TIM10 Antibody; Anti-TIM10A Antibody; anti-TIMM10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human TIMM10
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-TIMM10 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry)
Immunogen	E coli-derived Human TIMM10 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human TIMM10 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




IP (Immunoprecipitation)

TIMM10 was immunoprecipitated using: Lane A: 0.5 mg HepG2 Whole Cell Lysate, 4 uL anti-TIMM10 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-TIMM10 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 10 kDa. Observed band size: 10 kDa.



IHC (Immunohistochemistry)

Immunochemical staining of human TIMM10 in human liver with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human TIMM10 in human kidney with rabbit polyclonal antibody at 1:300 dilution, formalin-fixed paraffin embedded sections.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/timm10-antibody-immunoprecipitation-immunohistochemistry-elisa/259079>