

Rabbit anti-Human PTK2/FAK Polyclonal Antibody | anti-PTK2/FAK antibody

Anti-PTK2/FAK Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259092	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PTK2/FAK; Polyclonal Antibody; Anti-PTK2/FAK Antibody; Rabbit Polyclonal; protein tyrosine kinase 2; Anti-FADK Antibody; Anti-FAK Antibody; Anti-FAK1 Antibody; Anti-FRNK Antibody; Anti-p125FAK Antibody; Anti-pp125FAK Antibody; Anti-PPP1R71 Antibody; anti-PTK2/FAK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human PTK2/FAK
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-PTK2/FAK antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human PTK2/FAK1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human PTK2/FAK1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES




WB (Western Blot)

Anti-PTK2/FAK1 rabbit polyclonal antibody at 1:500 dilution Lane A: HeLa Whole Cell Lysate Lysates/proteins at 30 ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 119 kDa Observed band size: 119 kDa




IHC (Immunohistochemistry)

Immunochemical staining of human PTK2 in human testis with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human PTK2 in human kidney with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



IP (Immunoprecipitation)

PTK2 was immunoprecipitated using: Lane A: 0.5 mg HeLa Whole Cell Lysate 4 uL anti-PTK2 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-PTK2 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 119 kDa. Observed band size: 120 kDa.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ptk2-fak-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/259092>