

Rabbit anti-Human CHORDC1 Polyclonal Antibody | anti-CHORDC1 antibody

Anti-CHORDC1 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259093	Rabbit	Human	Immunoprecipitation, Immunofluorescence, Immunocytochemistry


PRODUCT SYNONYMS

CHORDC1; Polyclonal Antibody; Anti-CHORDC1 Antibody; Rabbit Polyclonal; cysteine and histidine-rich domain (CHORD) containing 1; Anti-CHP1 Antibody; anti-CHORDC1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CHORDC1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CHORDC1 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), ICC (Immunocytochemistry)
Immunogen	E coli-derived Human CHORDC1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human CHORDC1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IP (Immunoprecipitation)

CHORDC1 was immunoprecipitated using: Lane A: 0.5 mg 293 Whole Cell Lysate 4 uL anti-CHORDC1 rabbit polyclonal antibody and 60 ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CHORDC1 rabbit polyclonal antibody, at 1:100 dilution. Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 37 kDa. Observed band size: 37 kDa.



IF (Immunofluorescence)

Immunofluorescence staining of CHORDC1 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human CHORDC1 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor 488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/chordc1-antibody-immunoprecipitation-immunofluorescence-immunocytochemistry-elisa/259093>