

Rabbit anti-Human HUWE1 Polyclonal Antibody | anti-HUWE1 antibody

Anti-HUWE1 Antibody, Rabbit Polyclonal

CATALOG # AAA259110	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

HUWE1; Polyclonal Antibody; Anti-HUWE1 Antibody; Rabbit Polyclonal; anti-HUWE1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human HUWE1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-HUWE1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human HUWE1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human HUWE1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles.

IMAGES



IHC (Immunohistochemistry)

Immunochemical staining of human HUWE1 in human testis with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-HUWE1 rabbit polyclonal antibody at 1:500 dilution. Lane A: 293T Whole Cell Lysate. Lane B: HeLa Whole Cell Lysate. Lysates/proteins at 30 ug per lane. Secondary: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 481 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-HUWE1 antibody	HUWE1 (HECT, UBA And WWE Domain Containing E3 Ubiquitin Protein Ligase 1) is a large proteins. There are conflicting reports of the role of HUWE1 in regulating the oncogene c-MYC (MYC). In one model, HUWE1 regulates the stability of MYC through K48-linked ubiquitination and knockdown of HUWE1 leads to increased MYC and drives tumor progression [16]. Conversely, in other tumor types HUWE1 activates transcription of MYC through K63-linked ubiquitination and HUWE1 silencing or inhibition prevents MYC signaling and reduces tumor cell proliferation. HUWE1 may play a cell-type and context-dependent role in MYC regulation.
---	---

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/huwe1-antibody-immunohistochemistry-western-blot-elisa/259110>