

SARS-CoV-2 (2019-nCoV) Recombinant Protein | SARS-COV-2 recombinant protein

SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein, Biotinylated


CATALOG # AAA259356	HOST HEK293 Cells	REACTIVITY SARS-CoV-2
-------------------------------	-----------------------------	---------------------------------

PRODUCT SYNONYMS


SARS-CoV-2 (2019-nCoV); SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein; Biotinylated; SARS-COV-2 recombinant protein

PRODUCT SPECIFICATIONS

Host	HEK293 Cells
Reactivity	SARS-CoV-2
Purity/Purification	> 95% as determined by SDS-PAGE
Form/Format	Lyophilized from sterile PBS, pH 7.4. Normally 5%-8% trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Sequence	Arg319-Phe 541
Tag	C-His
Predicted N Terminal	Arg319
Endotoxin	< 1.0 EU perug protein as determined by the LAL method.
Protein Construction	A DNA sequence encoding the SARS-CoV-2 (2019-nCoV) Spike Protein (RBD)(YP_009724390.1)(Arg319-Phe541) was expressed with a polyhistidine tag at the C-terminus. The purified protein was biotinylated in vitro.
Activity	Measured by its binding ability in a functional ELISA. Immobilized human ACE2 protein (Fc tag) at 2ug/mL (100uL/well) can bind SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein, Biotinylated, the EC50 of SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein, Biotinylated is 20-60ng/mL.
Preparation and Storage	Samples are stable for up to twelve months from date of receipt at -20 degree C to -80 degree C. Store it under sterile conditions at -20 degree C to -80 degree C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles. In general, recombinant proteins are provided as lyophilized powder which are shipped at ambient temperature. Bulk packages of recombinant proteins are provided as frozen liquid. They are shipped out with blue ice unless customers require otherwise.



Application Data



Application Data
Measured by its binding ability in a functional ELISA. Immobilized human ACE2 protein (Fc tag)(10108-H05H) at 2ug/mL (100uL/well) can bind SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein, Biotinylated (AAA259356), the EC50 of SARS-CoV-2 (2019-nCoV) Spike RBD-His Recombinant Protein, Biotinylated (AAA259356) is 20-60ng/mL.

ADDITIONAL INFORMATION

Related Product Information for SARS-CoV-2 recombinant protein	The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell. Known receptors bind S1 are ACE2, angiotensin-converting enzyme 2; DPP4, dipeptidyl peptidase-4; APN, aminopeptidase N; CEACAM, carcinoembryonic antigen-related cell adhesion molecule 1; Sia, sialic acid; O-ac Sia, O-acetylated sialic acid. The spike is essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together. The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. It's been reported that SARS-CoV-2 (COVID-19 coronavirus, 2019-nCoV) can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity. The main functions for the Spike protein are summarized as: Mediate receptor binding and membrane fusion; Defines the range of the hosts and specificity of the virus; Main component to bind with the neutralizing antibody; Key target for vaccine design; Can be transmitted between different hosts through gene recombination or mutation of the receptor binding domain (RBD), leading to a higher mortality rate.
References	She. S. e. al. (2007. Expression. glycosylation. an. modificatio. o. th. spik. (S. glycoprotei. o. SAR. CoV. Method. Mo. Biol. 379. 127-35.D. L. e. al. (2009. Th. spik. protei. o. SARS-CoV--. targe. fo. vaccin. an. therapeuti. development. Na. Re. Microbiol. . (3). 226-36.Xia. X. e. al. (2004. Th. SARS-Co. . glycoprotein. Cel. Mo. Lif. Sci. 6. (19-20). 2428-30.

NCBI AND UNIPROT INFORMATION

Molecular Weight	51.5 kDa
------------------	----------

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/sars-cov-2-2019-ncov-recombinant-protein-hek293-cells/259356>