

Rabbit anti-Human RBM15 Polyclonal Antibody | anti-RBM15 antibody

Anti-RBM15 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259435	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RBM15; Polyclonal Antibody; Anti-RBM15 Antibody; Rabbit Polyclonal; RBM15 Antibody; Rabbit PAb; Antigen Affinity Purified; RNA binding motif protein 15; Anti-OTT Antibody; Anti-OTT1 Antibody; Anti-SPEN Antibody; anti-RBM15 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human RBM15
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-RBM15 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human RBM15 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human RBM15 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES



IHC (Immunohistochemistry)
Immunochemical staining of human RBM15 in human placenta with rabbit polyclonal antibody at 1:100 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)
Anti-RBM15 rabbit polyclonal antibody at 1:500 dilutionLane A: HeLa Whole Cell LysateLane B: A431 Whole Cell LysateLysates/proteins at 30ug per lane.SecondaryGoat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.Developed using the ECL technique. Performed under reducing conditions.Predicted band size:107 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-RBM15 antibody

RBM15, an RNA-binding protein, plays important roles in the growth and apoptosis of cells, especially blood cells through regulating multiple signal pathways such as Notch and Wnt. An increasing body of evidence has suggested that RBM15 may play a key function on the development of various blood diseases, such as acute/chronic myeloid leukemia and kaposi's sarcoma. Overexpression of PRMT1 in acute megakaryocytic leukemia cell lines blocks megakaryocyte terminal differentiation by downregulation of RBM15 protein level. Gene silencing of RBM15 leads to cytoplasmic depletion and nuclear accumulation of general mRNA as well as individual endogenous transcripts, indicating that RBM15 is required for efficient mRNA export.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/rbm15-antibody-immunohistochemistry-western-blot-elisa/259435>