

Rabbit anti-Human CARD11 Polyclonal Antibody | anti-CARD11 antibody

Anti-CARD11 Antibody, Rabbit Polyclonal

CATALOG # AAA259439	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

CARD11; Polyclonal Antibody; Anti-CARD11 Antibody; Rabbit Polyclonal; CARD11 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-BENTA Antibody; Anti-BIMP3 Antibody; Anti-CARMA1 Antibody; Anti-IMD11 Antibody; Anti-IMD11A Antibody; Anti-PPBL Antibody; anti-CARD11 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CARD11
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CARD11 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human CARD11 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human CARD11 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES




IHC (Immunohistochemistry)

Immunochemical staining of human CARD11 in human urinary bladder with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



IHC (Immunohistochemistry)

Immunochemical staining of human CARD11 in human kidney with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



WB (Western Blot)

Anti-CARD11 rabbit polyclonal antibody at 1:500 dilution Lane A: Jurkat Whole Cell Lysate Lysates/proteins at 30ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 133 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/card11-antibody-immunohistochemistry-western-blot-elisa/259439>