

Rabbit anti-Human CCAR1 Polyclonal Antibody | anti-CCAR1 antibody

Anti-CCAR1 Antibody, Rabbit Polyclonal

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259453	Rabbit	Human	Immunoprecipitation, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CCAR1; Polyclonal Antibody; Anti-CCAR1 Antibody; Rabbit Polyclonal; CCAR1 Antibody; Rabbit PAb; Antigen Affinity Purified; Anti-CARP-1 Antibody; Anti-CARP1 Antibody; Anti-DIS Antibody; Anti-FLJ10590 Antibody; anti-CCAR1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Human CCAR1
Purity/Purification	Protein A & Antigen Affinity
Form/Format	Liquid; PBS, pH7.0 with 0.03% Proclin300
Applicable Applications for anti-CCAR1 antibody	IP (Immunoprecipitation), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	E coli-derived Human CCAR1 fragment
Preparation	Produced in rabbits immunized with E coli-derived Human CCAR1 fragment, and purified by antigen affinity chromatography.
Preparation and Storage	This antibody can be stored at 2 degree C-8 degree C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20 degree C to -80 degree C. Avoid repeated freeze-thaw cycles. This antibody is shipped as liquid solution at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

IMAGES



IP (Immunoprecipitation)

CCAR1 was immunoprecipitated using: Lane A:0.5 mg HeLa Whole Cell Lysate Lane B:0.5 mg Jurkat Whole Cell Lysate 4 uL anti-CCAR1 rabbit polyclonal antibody and 60ug of Immunomagnetic beads Protein A/G. Primary antibody: Anti-CCAR1 rabbit polyclonal antibody, at 1:100 dilution Secondary antibody: Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 132 kDa Observed band size :150 kDa




IHC (Immunohistochemistry)

Immunochemical staining of human CCAR1 in human liver with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.

**IHC (Immunohistochemistry)**

Immunochemical staining of human CCAR1 in human lymph node with rabbit polyclonal antibody at 1:2000 dilution, formalin-fixed paraffin embedded sections.

**WB (Western Blot)**

Anti-CCAR1 rabbit polyclonal antibody at 1:2000 dilution Lane A: HeLa Whole Cell Lysate Lane B: Jurkat Whole Cell Lysate Lysates/proteins at 30ug per lane. Secondary Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution. Developed using the ECL technique. Performed under reducing conditions. Predicted band size: 132 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/ccar1-antibody-immunoprecipitation-immunohistochemistry-western-blot-elisa/259453>