

Rabbit anti-Human EG5 Polyclonal Antibody | anti-KIF11 antibody

Anti-EG5 (pT926) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259600	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

EG5; Polyclonal Antibody; Anti-EG5 (pT926) Phospho Antibody; KNSL1; TRIP5; Kinesin-like protein KIF11; Kinesin-like protein 1; Kinesin-like spindle protein HKSP; Kinesin-related motor protein Eg5; Thyroid receptor-interacting protein 5; TR-interacting protein 5; TRIP-5; anti-KIF11 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of EG5 (pT926) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1056
Applicable Applications for anti-KIF11 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human EG5 (pT926). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of EG5 (pT926) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of EG5 (pT926) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of EG5 (pT926) expression in Jurkat (A), HEK293T (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KIF11 antibody	Rabbit polyclonal antibody to EG5 (pT926)
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	13699824
NCBI GeneID	3832
NCBI Accession #	NP_004514.2
NCBI GenBank Nucleotide #	NM_004523.3
UniProt Accession #	P52732; Q15716; Q5VWX0; A0AV49; B2RMV3
Molecular Weight	119,159 Da
NCBI Official Full Name	kinesin-like protein KIF11
NCBI Official Synonym Full Names	kinesin family member 11
NCBI Official Symbol	KIF11
NCBI Official Synonym Symbols	EG5; HKSP; KNSL1; MCLMR; TRIP5
NCBI Protein Information	kinesin-like protein KIF11
UniProt Protein Name	Kinesin-like protein KIF11
UniProt Gene Name	KIF11
UniProt Synonym Gene Names	EG5; KNSL1; TRIP5; TR-interacting protein 5; TRIP-5
UniProt Entry Name	KIF11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/eg5-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259600>