

Rabbit anti-Human p47 phox Polyclonal Antibody | anti-NCF1 antibody

Anti-p47 phox (pS359) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259638	Rabbit	Human	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

p47 phox; Polyclonal Antibody; Anti-p47 phox (pS359) Phospho Antibody; NOXO2; SH3PXD1A; Neutrophil cytosol factor 1; NCF-1; 47 kDa autosomal chronic granulomatous disease protein; 47 kDa neutrophil oxidase factor; NCF-47K; Neutrophil NADPH oxidase factor 1; Nox organizer 2; Nox-organizing protein 2; SH3 and PX domain-containing protein 1A; p47-phox; anti-NCF1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of p47 phox (pS359) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	390
Applicable Applications for anti-NCF1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human p47 phox (pS359). The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of p47 phox (pS359) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of p47 phox (pS359) staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of p47 phox (pS359) expression in HeLa (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-NCF1 antibody	Rabbit polyclonal antibody to p47 phox (pS359)
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	528078313
NCBI GeneID	653361
NCBI Accession #	NP_000256.4
NCBI GenBank Nucleotide #	NM_000265.5
UniProt Accession #	P14598; O43842; Q2PP07; Q53FR5; Q9BU90; Q9BXI7; Q9BXI8; Q9UDV9; Q9UMU2; A6NEH2; A8K7S9
Molecular Weight	30,781 Da
NCBI Official Full Name	neutrophil cytosol factor 1
NCBI Official Synonym Full Names	neutrophil cytosolic factor 1
NCBI Official Symbol	NCF1
NCBI Official Synonym Symbols	NCF1A; NOXO2; p47phox; SH3PXD1A
NCBI Protein Information	neutrophil cytosol factor 1
UniProt Protein Name	Neutrophil cytosol factor 1
UniProt Gene Name	NCF1
UniProt Synonym Gene Names	NOXO2; SH3PXD1A; NCF-1
UniProt Entry Name	NCF1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.