

Rabbit Beta-catenin Polyclonal Antibody | anti-CTNNB1 antibody

Anti-Beta-catenin (pS33) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259650	Rabbit	Human, Mouse, Rat, Bovine, Dog, Porcine, Rabbit, Sheep	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Beta-catenin; Polyclonal Antibody; Anti-Beta-catenin (pS33) Phospho Antibody; CTNNB; Catenin beta-1; anti-CTNNB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Porcine, Rabbit, Sheep
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Beta-catenin (pS33) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	88
Applicable Applications for anti-CTNNB1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Beta-catenin (pS33). The exact sequence is proprietary.
Entrez Gene	1499 (Human); 12387 (Mouse); 84353 (Rat)
SwissProt	P35222 (Human); Q02248 (Mouse); Q9WU82 (Rat)
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Beta-catenin (pS33) staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Beta-catenin (pS33) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Beta-catenin (pS33) expression in HeLa (A), HepG2 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CTNNB1 antibody	Rabbit polyclonal antibody to Beta-catenin (pS33)
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	148233338
NCBI GeneID	1499
NCBI Accession #	NP_001091679.1
NCBI GenBank Nucleotide #	NM_001098209.1
UniProt Accession #	P35222; Q8NEW9; Q8NI94; Q9H391; A8K1L7
Molecular Weight	9,501 Da
NCBI Official Full Name	catenin beta-1
NCBI Official Synonym Full Names	catenin (cadherin-associated protein), beta 1, 88kDa
NCBI Official Symbol	CTNNB1

NCBI Official Synonym Symbols	CTNNB; MRD19; armadillo
NCBI Protein Information	catenin beta-1
UniProt Protein Name	Catenin beta-1
UniProt Gene Name	CTNNB1
UniProt Synonym Gene Names	CTNNB
UniProt Entry Name	CTNB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/beta-catenin-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259650>