

Rabbit ERCC5 Polyclonal Antibody | anti-ERCC5 antibody

Anti-ERCC5 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259697	Rabbit	Human, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

ERCC5; Polyclonal Antibody; Anti-ERCC5 Antibody; ERCM2; XPG; XPGC; DNA repair protein complementing XP-G cells; DNA excision repair protein ERCC-5; Xeroderma pigmentosum group G-complementing protein; anti-ERCC5 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of ERCC5 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1186
Applicable Applications for anti-ERCC5 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ERCC5. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ERCC5 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of ERCC5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ERCC5 expression in COLO205 (A), GM00637 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ERCC5 antibody	Rabbit polyclonal antibody to ERCC5
---	-------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	51988900
NCBI GeneID	2073
NCBI Accession #	NP_000114.2
NCBI GenBank Nucleotide #	NM_000123.3
UniProt Accession #	P28715; Q5JUS4; Q5JUS5; Q7Z2V3; Q8IZL6; Q8N1B7; Q9HD59; Q9HD60; A6NGT4
Molecular Weight	27,259 Da
NCBI Official Full Name	DNA repair protein complementing XP-G cells
NCBI Official Synonym Full Names	excision repair cross-complementation group 5
NCBI Official Symbol	ERCC5
NCBI Official Synonym Symbols	XPG; UVDR; XPGC; COFS3; ERCM2; ERCC5-201
NCBI Protein Information	DNA repair protein complementing XP-G cells
UniProt Protein Name	DNA repair protein complementing XP-G cells
UniProt Gene Name	ERCC5
UniProt Synonym Gene Names	ERCM2; XPG; XPGC
UniProt Entry Name	ERCC5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ercc5-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259697>