

Rabbit Caspase 6 Polyclonal Antibody | anti-CASP6 antibody

Anti-Caspase 6 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259704	Rabbit	Human, Mouse, Rat, Bovine, Monkey	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Caspase 6; Polyclonal Antibody; Anti-Caspase 6 Antibody; MCH2; Caspase-6; CASP-6; Apoptotic protease Mch-2; anti-CASP6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Monkey
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Caspase 6 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	293
Applicable Applications for anti-CASP6 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Caspase 6. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Caspase 6 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Caspase 6 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Caspase 6 expression in HepG2 (A), Raw264.7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-CASP6 antibody	Rabbit polyclonal antibody to Caspase 6
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	14916483
NCBI GeneID	839
NCBI Accession #	NP_001217.2
NCBI GenBank Nucleotide #	NM_001226.3
UniProt Accession #	P55212; Q9BQE7
Molecular Weight	22,574 Da
NCBI Official Full Name	caspase-6 isoform alpha preproprotein
NCBI Official Synonym Full Names	caspase 6, apoptosis-related cysteine peptidase
NCBI Official Symbol	CASP6
NCBI Official Synonym Symbols	MCH2
NCBI Protein Information	caspase-6
UniProt Protein Name	Caspase-6
UniProt Gene Name	CASP6
UniProt Synonym Gene Names	MCH2; CASP-6
UniProt Entry Name	CASP6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/caspase-6-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259704>