

Rabbit Alpha-adducin Polyclonal Antibody | anti-ADD1 antibody

Anti-Alpha-adducin Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259712	Rabbit	Human, Mouse, Rat, Bovine	Immunoprecipitation, Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Alpha-adducin; Polyclonal Antibody; Anti-Alpha-adducin Antibody; ADD1; ADDA; Erythrocyte adducin subunit alpha; ADD2; ADDB; Beta-adducin; Erythrocyte adducin subunit beta; anti-ADD1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Alpha-adducin protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	737
Applicable Applications for anti-ADD1 antibody	IP (Immunoprecipitation), ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Alpha-adducin. The exact sequence is proprietary.
Preparation and Storage	Upon receipt aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Alpha-adducin staining in HEK293 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Alpha-adducin staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Alpha-adducin expression in Jurkat (A), HeLa (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ADD1 antibody	Rabbit polyclonal antibody to Alpha-adducin
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	29826319
NCBI GeneID	118
NCBI Accession #	NP_001110.2
NCBI GenBank Nucleotide #	NM_001119.4
UniProt Accession #	P35611; Q13734; Q14729; Q16156; Q86XM2; Q9UJB6; A2A3N8; A2A3P0; B4DI79; D3DVR3; D3DVR4; D3DVR5
NCBI Official Full Name	alpha-adducin isoform a
NCBI Official Synonym Full Names	adducin 1 (alpha)
NCBI Official Symbol	ADD1
NCBI Official Synonym Symbols	ADDA
NCBI Protein Information	alpha-adducin
UniProt Protein Name	Alpha-adducin
UniProt Gene Name	ADD1
UniProt Synonym Gene Names	ADDA
UniProt Entry Name	ADDA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/alpha-adducin-antibody-immunoprecipitation-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259712>