

HER4 Polyclonal Antibody | anti-ERBB4 antibody

Anti-HER4 Antibody

CATALOG # AAA259724	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	--	--


PRODUCT SYNONYMS

HER4; Polyclonal Antibody; Anti-HER4 Antibody; Receptor tyrosine-protein kinase erbB-4; Proto-oncogene-like protein c-ErbB-4; Tyrosine kinase-type cell surface receptor HER4; p180erbB4; anti-ERBB4 antibody

PRODUCT SPECIFICATIONS


Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of HER4 protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1292
Applicable Applications for anti-ERBB4 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Source	Rabbit
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human HER4. The exact sequence is proprietary.
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of HER4 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of HER4 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of HER4 expression in Y79 (A), MOLT4 (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ERBB4 antibody	Rabbit polyclonal antibody to HER4
---	------------------------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	110825958
NCBI GeneID	2066
NCBI Accession #	NP_001036064.1
NCBI GenBank Nucleotide #	NM_001042599.1
UniProt Accession #	Q15303; Q2M1W1; Q59EW4; B7ZLD7; B7ZLE2; B7ZLE3
NCBI Official Full Name	receptor tyrosine-protein kinase erbB-4 isoform JM-a/CVT-2
NCBI Official Synonym Full Names	erb-b2 receptor tyrosine kinase 4
NCBI Official Symbol	ERBB4
NCBI Official Synonym Symbols	HER4; ALS19; p180erbB4
NCBI Protein Information	receptor tyrosine-protein kinase erbB-4
UniProt Protein Name	Receptor tyrosine-protein kinase erbB-4
UniProt Gene Name	ERBB4
UniProt Synonym Gene Names	HER4; 4ICD; E4ICD
UniProt Entry Name	ERBB4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.