

Rabbit VEGFR2 Polyclonal Antibody | anti-KDR antibody

Anti-VEGFR2 (pY1175) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259744	Rabbit	Human, Mouse, Rat, Pig, Sheep	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

VEGFR2; Polyclonal Antibody; Anti-VEGFR2 (pY1175) Phospho Antibody; FLK1; Vascular endothelial growth factor receptor 2; VEGFR-2; Fetal liver kinase 1; FLK-1; Kinase insert domain receptor; KDR; Protein-tyrosine kinase receptor flk-1; CD309; anti-KDR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Pig, Sheep
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of VEGFR2 (pY1175) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	1356
Applicable Applications for anti-KDR antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human VEGFR2 (pY1175). The exact sequence is proprietary.
Preparation and Storage	Upon receipt aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of VEGFR2 (pY1175) staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of VEGFR2 (pY1175) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of VEGFR2 (pY1175) expression in HEK293T EGF-treated (A), NIH3T3 EGF-treated (B) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-KDR antibody

Rabbit polyclonal antibody to VEGFR2 (pY1175)

NCBI AND UNIPROT INFORMATION

NCBI GI #	11321597
NCBI GeneID	3791
NCBI Accession #	NP_002244.1
NCBI GenBank Nucleotide #	NM_002253.2
UniProt Accession #	P35968; O60723; Q14178; A2RRS0; B5A925; C5IFA0
NCBI Official Full Name	vascular endothelial growth factor receptor 2
NCBI Official Synonym Full Names	kinase insert domain receptor
NCBI Official Symbol	KDR
NCBI Official Synonym Symbols	FLK1; CD309; VEGFR; VEGFR2
NCBI Protein Information	vascular endothelial growth factor receptor 2
UniProt Protein Name	Vascular endothelial growth factor receptor 2
UniProt Gene Name	KDR
UniProt Synonym Gene Names	FLK1; VEGFR2; VEGFR-2; FLK-1; KDR
UniProt Entry Name	VGFR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/vegfr2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/259744>