

Rabbit Estrogen Receptor alpha Polyclonal Antibody | anti-ESR1 antibody

Anti-Estrogen Receptor alpha (pS106) Phospho Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA259780	Rabbit	Human, Mouse, Rat, Monkey, Rabbit	Immunocytochemistry, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Estrogen Receptor alpha; Polyclonal Antibody; Anti-Estrogen Receptor alpha (pS106) Phospho Antibody; ESR; NR3A1; Estrogen receptor; ER; ER-alpha; Estradiol receptor; Nuclear receptor subfamily 3 group A member 1; anti-ESR1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey, Rabbit
Clonality	Polyclonal
Specificity	Recognizes endogenous levels of Estrogen Receptor alpha (pS106) protein.
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Sequence Length	595
Applicable Applications for anti-ESR1 antibody	ICC (Immunocytochemistry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Estrogen Receptor alpha (pS106). The exact sequence is proprietary.
Preparation and Storage	Upon delivery aliquot and store at -20 degree C for one year. Avoid freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Estrogen Receptor alpha (pS106) staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Estrogen Receptor alpha (pS106) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
 Western blot analysis of Estrogen Receptor alpha (pS106) expression in MCF7 EGF-treated
 (A) whole cell lysates.

ADDITIONAL INFORMATION

Related Product Information for anti-ESR1 antibody	Rabbit polyclonal antibody to Estrogen Receptor alpha (pS106)
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	62821794
NCBI GeneID	2099
NCBI Accession #	NP_000116.2
NCBI GenBank Nucleotide #	NM_000125.3
UniProt Accession #	P03372; Q13511; Q14276; Q5T5H7; Q6MZQ9; Q9NU51; Q9UDZ7; Q9UIS7
NCBI Official Full Name	estrogen receptor isoform 1
NCBI Official Synonym Full Names	estrogen receptor 1
NCBI Official Symbol	ESR1
NCBI Official Synonym Symbols	ER; ESR; Era; ESRA; ESTRR; NR3A1
NCBI Protein Information	estrogen receptor
UniProt Protein Name	Estrogen receptor
UniProt Gene Name	ESR1
UniProt Synonym Gene Names	ESR; NR3A1; ER
UniProt Entry Name	ESR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/estrogen-receptor-alpha-antibody-immunocytochemistry-immunofluorescence-immunohistochemistry-western-blot-elisa/259780>